



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Correlations between fMRI activation and individual psychotic symptoms in un-medicated subjects at high genetic risk of schizophrenia

Citation for published version:

Whalley, HC, Gountouna, V-E, Hall, J, McIntosh, A, Whyte, M-C, Simonotto, E, Job, DE, Owens, DGC, Johnstone, EC & Lawrie, SM 2007, 'Correlations between fMRI activation and individual psychotic symptoms in un-medicated subjects at high genetic risk of schizophrenia', *BMC Psychiatry*, vol. 7, 61. <https://doi.org/10.1186/1471-244X-7-61>

Digital Object Identifier (DOI):

[10.1186/1471-244X-7-61](https://doi.org/10.1186/1471-244X-7-61)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

BMC Psychiatry

Publisher Rights Statement:

© 2007 Whalley et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Research article

Open Access

Correlations between fMRI activation and individual psychotic symptoms in un-medicated subjects at high genetic risk of schizophrenia

Heather C Whalley*, Viktoria-Eleni Gountouna, Jeremy Hall, Andrew McIntosh, Marie-Claire Whyte, Enrico Simonotto, Dominic E Job, David GC Owens, Eve C Johnstone and Stephen M Lawrie

Address: Division of Psychiatry, School of Molecular and Clinical Medicine, University of Edinburgh, Edinburgh, Scotland, UK.

Email: Heather C Whalley* - hwhalley@staffmail.ed.ac.uk; Viktoria-Eleni Gountouna - E.Gountouna@ed.ac.uk; Jeremy Hall - jhall5@staffmail.ed.ac.uk; Andrew McIntosh - andrew.mcintosh@ed.ac.uk; Marie-Claire Whyte - marieclaire.mackie@gmail.com; Enrico Simonotto - esimonotto@invivocorp.com; Dominic E Job - djob@staffmail.ed.ac.uk; David GC Owens - david.owens@ed.ac.uk; Eve C Johnstone - e.johnstone@ed.ac.uk; Stephen M Lawrie - s.lawrie@ed.ac.uk

* Corresponding author

Published: 29 October 2007

Received: 11 May 2007

BMC Psychiatry 2007, 7:61 doi:10.1186/1471-244X-7-61

Accepted: 29 October 2007

This article is available from: <http://www.biomedcentral.com/1471-244X/7/61>

© 2007 Whalley et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: It has been proposed that different types of psychopathology in schizophrenia may reflect distinguishable pathological processes. In the current study we aimed to address such associations in the absence of confounders such as medication and disease chronicity by examining specific relationships between fMRI activation and individual symptom severity scores in un-medicated subjects at high genetic risk of schizophrenia.

Methods: Associations were examined across two functional imaging paradigms: the Hayling sentence completion task, and an encoding/retrieval task, comprising encoding (at word classification) and retrieval (old word/new word judgement). Symptom severity was assessed using the positive and negative syndrome scale (PANSS). Items examined were hallucinations, delusions, and suspiciousness/persecution.

Results: Associations were seen in the anterior middle temporal gyrus in relation to hallucination scores during the sentence completion task, and in the medial temporal lobe in association with suspiciousness/persecution scores in the encoding/retrieval task. Cerebellar activation was associated with delusions and suspiciousness/persecution scores across both tasks with differing patterns of laterality.

Conclusion: These results support a role for the lateral temporal cortex in hallucinations and medial temporal lobe in positive psychotic symptoms. They also highlight the potential role of the cerebellum in the formation of delusions. That the current results are seen in un-medicated high risk subjects indicates these associations are not specific to the established illness and are not related to medication effects.

Background

Schizophrenia is characterised by the presence of positive psychotic symptoms, particularly hallucinations and delusions. The form and content of these symptoms can vary considerably between patients and it has been suggested that different types of psychopathology may reflect distinguishable pathological processes [1]. A number of approaches have been taken to examine such associations [2], including scanning patients with specific symptoms against a suitable control group, or examining patients during the active experience of individual symptoms. Neural substrates implicated in positive psychotic symptoms have included medial temporal lobe (MTL) structures [3-5], lateral temporal cortex [6], and inferior frontal cortex [6-8].

Those studies that have further split positive psychotic symptoms have implicated the lateral temporal cortex in auditory hallucinations, primarily auditory and language related areas in the superior temporal gyrus [9-15]. However, other regions have also been reported to be associated with auditory hallucinations including medial and anterior temporal lobe [10,13,14,16], medial and lateral prefrontal regions [10-12,14,16,17], and cerebellum [12,13]. Far fewer studies have examined the neural correlates of delusions and results are at present inconclusive [18-20].

Such studies are however often small and confounded by antipsychotic medication effects. An alternative approach is to examine a larger group of subjects who are perhaps less severely affected, better able to co-operate with testing, and less likely to be prescribed antipsychotic medication. Individual psychotic symptoms have been noted in the general population (~10%) [21], and have been shown to be more prevalent in high risk samples (~40%) [22]. In the Edinburgh High Risk Study psychotic symptoms were associated with, although not necessarily followed by, subsequent conversion to schizophrenia (as only about 25% converted). Examining relationships between brain activation and the severity of sub-diagnostic psychotic symptoms in high risk subjects in the absence of anti-psychotic medication effects may improve our understanding of the biological basis of the early stages of schizophrenia.

In the current study our aim was to map the neural underpinnings of hallucinations and delusions. Current neuropsychological models of these symptoms stress the importance of disruptions in language processing and mnemonic functions [1,23]. The tasks used in the current study (a verbal fluency task and a verbal encoding/retrieval task) were chosen on the basis of their involvement in these networks and cognitive processes. Based on the schizophrenia literature we hypothesised significant

correlations between the severity of hallucinations and delusions and activation in MTL structures, and between hallucination scores and activation in lateral temporal regions.

Methods

Subject details

The Edinburgh High Risk Study (EHRS) examined young adults aged 16–24 years at ascertainment (1995–1999) at enhanced genetic risk of schizophrenia (with two or more affected first or second degree relatives) over the period at which they are at greatest risk of becoming ill, in comparison with matched healthy controls with no family history of psychotic illness [22,24,25]. All subjects were antipsychotic naïve throughout the study. This report presents results obtained during the second phase of the study (1999–2004) when fMRI was introduced into the protocol. All subjects were supplied with detailed written information regarding the study and provided written informed consent. The study was approved by the Psychiatry and Clinical Psychology subcommittee of the Lothian research ethics committee. Since this was a within group correlation study the healthy control data were not included, further details are available in earlier publications [26,27].

A total of 96 high risk subjects underwent an fMRI scan which included two functional imaging paradigms; the Hayling sentence completion test (experiment 1) and the encoding/retrieval task (experiment 2). Three subjects were excluded from further analysis due to minor vascular malformations and a temporal lobe cyst. For experiment 1 a further four subjects were excluded due to movement artefact leaving 89 fMRI scans. For experiment 2, seven scans were excluded (five due to excessive movement and two due to loss of behavioural data) leaving 86 fMRI scans. These samples included 69 high risk subjects previously reported [26,27], with an additional 20 subjects representing the completed cohort. There was an overlap of 83 subjects who provided usable scans for both paradigms. For the sentence completion task the mean age was 26.2 years (s.d. 3.3), comprising 42 males and 47 females. For the encoding/retrieval task the mean age was 26.1 years (s.d. 3.3), comprising 41 males and 45 females. The majority of subjects were working or in full time higher education [22]. None of the subjects were on anti-psychotic medication, seeking treatment or saw themselves as unwell, nor did they fulfil diagnostic criteria for any psychiatric disorder.

The severity of positive psychotic symptoms was determined using the positive and negative syndrome scale (PANSS) [28] conducted by two experienced clinicians who had previously undergone the PANSS training programme and had reached the set standard of reliability

[29]. Since the primary interest of the current report were activations associated with delusions and hallucinations, the PANSS scores for these measures (first, third, and sixth items on the positive scale representing delusions, hallucinations and suspiciousness/persecution) were entered into further analysis. The score for suspiciousness/persecution was selected along with the score for delusions as it is considered to reflect a paranoid element.

Scanning procedure

Imaging was carried out at the Brain Imaging Research Centre (BIRC) for Scotland on a GE 1.5 T Signa scanner (GE Medical, Milwaukee, USA). The imaging protocol consisted of a localiser scan, followed by a T2-weighted fast spin-echo sequence, and a structural T1 weighted sequence followed by the functional imaging paradigms. For experiment 1 axial gradient-echo planar images (EPI) (TR/TE = 4000/40 ms; matrix = 64×128 ; field of view (fov) 220×440 mm) were acquired continually. Thirty eight contiguous 5-mm slices aligned to the anterior and posterior commissure were acquired within each TR period. Each acquisition was run for 204 volumes, of which the first four volumes were discarded. For experiment 2 functional images were again collected using an EPI sequence (TR/TE = 2000/40 ms; matrix = 64×64 ; field of view (fov) 220×220 mm). Twenty-four contiguous 5 mm axial (horizontal) slices were collected at an oblique angle aligned with the anterior and posterior commissure. Data were acquired during 2 sessions, consisting of 104 volumes for the first (word classification) and 204 volumes for the second session (word retrieval), for each session the first four volumes were discarded.

Experiment 1: Hayling sentence completion task (blocked design)

Participants performed the verbal initiation section of the Hayling sentence completion test in the scanner as described previously [26]. Subjects were shown sentences with the last word missing and were asked to silently think of an appropriate word to complete the sentence and press a button when they had done so. Sentences were presented in blocks of fixed difficulty, determined by the range of suitable words suggested by the sentence context, of which there were four levels. Each block lasted 40 seconds and included 8 sentences. Sentences were presented for a period of 3 seconds followed by a fixation cross for 2 seconds and subjects could respond at any time until the next sentence appeared. The rest condition consisted of viewing a screen of white circles on a black background for 40 seconds. The order of the blocks was pseudo-random, and each block was repeated four times. Immediately after scanning, subjects were given the same sequence of sentences on paper and requested to complete each sentence with the word they first thought of in the scanner. 'Word

appropriateness' scores were determined from the word frequency list of sentence completion norms [30].

Experiment 2: verbal encoding and retrieval (event-related)

During the word classification, or 'encoding', part of the task single words were presented in the scanner and subjects were asked to classify them as 'living' or 'non-living' by pressing a button. A total of 36 words were presented, 18 referring to living things and 18 to non-living things. During the 'retrieval' part of the task single words were presented randomly selected from those shown during the word classification condition along with matched similar new words and subjects were requested to classify them as either 'new' or 'old' words and signify their response by pressing a button. In total 72 words were presented, 36 words which were presented previously (old) intermixed with another 36 matched new words. For more detail see previous study [27]. Stimuli were presented for 2 seconds, followed by a variable fixation period of 2–10 seconds. Responses could be made at any time during presentation and subsequent fixation period by pressing a button. Both parts of the task were preceded by a practice session with feedback.

Image processing and analysis

Scan analysis was performed using the standard SPM approach in SPM2 [31] running in Matlab (The Math-Works, Natick, MA, USA). Briefly, EPI volumes were realigned to the mean volume in the series, normalised to a study specific EPI template (comprising 121 participants of the EHRS) and spatially smoothed with a $6 \times 6 \times 6$ mm³ FWHM Gaussian filter.

First level analysis

Statistical analysis was performed using the general linear model approach as implemented in SPM. Estimates of head movement from the realignment stage of pre-processing were included as additional regressors in the first level analysis. For experiment 1 contrasts were constructed to examine all four sentence completion conditions versus rest, and a parametric contrast which examines increasing activation with increasing difficulty. For the word classification condition of experiment 2 all items were treated as equal and entered into a single regressor model. One contrast of interest was generated examining word classification versus baseline experimental activation. For the retrieval condition three regressors were modelled, correct old words (correct recognition), correct new words (correct rejection) and incorrect words (further splitting of incorrect old and incorrect new was not considered appropriate due to insufficient events in each category). For the purposes of this study contrast images were constructed to examine correct recognition

versus baseline experimental activation and correct rejection versus baseline experimental activation.

Second level analysis

First level contrast images for each subject were then entered into a second level multiple regression analyses. For each of the five contrasts of interest a single model was used to examine correlations with the selected scores from the PANSS. The three symptom dimensions were modelled simultaneously within one design matrix hence variability common to two or more symptom dimensions is not reflected in the correlation values. In other words, the analysis performed represents, for example, the relationship between delusion severity and fMRI activation that is not accounted for by either suspiciousness or hallucination severity.

Statistical maps were thresholded at a level of $p = 0.005$ uncorrected, and regions were considered significant at $p < 0.05$ cluster level corrected for multiple comparisons as performed in SPM. Where significant relationships were found, data were extracted for each cluster (at the maximum voxel) and entered into SPSS (version 12) where a linear regression analysis was performed to check that the standardised residuals were normally distributed and to determine correlation coefficients between the individual symptom severity score and the degree of activation. All p values quoted in the text are at the corrected cluster level as performed in SPM. We would like to state however that positive and negative correlations were examined for each of the five contrasts separately (hence uncorrected voxel-wise p values are one-tailed) and correction for the number of these comparison was not performed. Since we had prior hypotheses regarding temporal lobe involvement in hallucinations we imposed an anatomically defined bilateral temporal lobe small volume correction (SVC) for the examination of correlations with the hallucinations measure [32]. Co-ordinates were converted from MNI (Montreal Neurological Institute) to Talairach co-ordinates using a non-linear transformation [33].

Results

Symptom scores and behavioural measures

The scores for all positive psychotic symptoms as measured by the PANSS are described in Table 1. The distribution of the three symptom dimensions examined in the current study are presented in Figure 1. Pair-wise correlations between the symptom scores are detailed in the figure legend. Behavioural measures (Table 2) indicated subjects were performing both tasks appropriately in the scanner and there were no significant correlations between these measures and any of three symptom severity scores. Previous analysis comparing high risk subjects to normal controls indicated no significant impairment in performance for either of the two tasks [26,27].

Group activation maps

For both experiments the within group maps for all contrasts of interest are shown in a Figure 2. These maps indicated regions commonly activated in these types of tasks and are consistent with previous results [26,27]. For sentence completion versus rest the main areas of activation were the left precentral gyrus, inferior frontal gyrus, medial/superior frontal gyrus, middle/superior temporal gyrus, cerebellum and occipital lobes bilaterally. The parametric contrast indicated increasing activation with increasing difficulty in left superior/medial frontal gyrus, inferior frontal gyrus and cerebellum.

For the encoding/retrieval task, word classification versus baseline experimental activation revealed large areas of activation encompassing bilateral inferior frontal gyrus, superior/medial frontal gyrus, insula, parietal lobe and cerebellum, along with basal ganglia, thalamus and hippocampus. Correct recognition and correct rejection versus experimental baseline revealed similar areas of activation including lateral and superior/medial frontal gyri, parietal lobe, insula, basal ganglia, thalamus and cerebellum. Activation was also observed in the hippocampus at its superior border although this formed part of a larger cluster encompassing basal ganglia and thalamus.

Table 1: PANSS measures for high risk subjects

| | Sentence completion (n = 89) | | Encoding/retrieval (n = 86) | |
|-------------------------------|------------------------------|-----------------------------------|-----------------------------|-----------------------------------|
| | Mean (SD) | Number of subjects with score > 2 | Mean (SD) | Number of subjects with score > 2 |
| P1 delusions | 1.51 (0.85) | 27 | 1.52 (0.86) | 27 |
| P2 conceptual disorganisation | 1.10 (0.40) | 6 | 1.10 (0.41) | 6 |
| P3 hallucinations | 1.31 (0.67) | 18 | 1.31 (0.67) | 17 |
| P4 excitement | 1.07 (0.30) | 5 | 1.07 (0.30) | 5 |
| P5 grandiosity | 1.07 (0.25) | 6 | 1.07 (0.26) | 6 |
| P6 suspiciousness/persecution | 1.28 (0.71) | 15 | 1.29 (0.72) | 15 |
| P7 hostility | 1.08 (0.35) | 5 | 1.08 (0.35) | 5 |
| PANSS positive total | 8.39 (2.4) | - | 8.43 (2.4) | - |




Figure 1
Distribution of symptom dimensions in high risk individuals. Values represent the number of subjects with PANSS scores >2 for each symptom type examined. Pair-wise correlation values between the symptom dimensions were as follows: Delusions × hallucinations $r = 0.34$ ($p < 0.01$); delusions × suspiciousness/persecution $r = 0.48$ ($p < 0.01$); hallucinations × suspiciousness/persecution $r = 0.32$ ($p < 0.01$).

Correlation with symptom severity

Experiment 1: Hayling sentence completion task

For sentence completion versus rest there was a positive correlation between activation in the left anterior middle temporal gyrus and the hallucinations score ($p = 0.01$, small volume correction for temporal lobes), see Figure 3a, Table 3.

Table 2: Behavioural measures

| Hayling sentence completion task | | |
|--|----------------------------------|--------------------------------|
| | Word appropriateness Mean (s.d.) | Reaction time (ms) Mean (s.d.) |
| Low constraint | 6.4 (0.97) | 2526 (597) |
| Medium low constraint | 3.3 (0.58) | 2427 (621) |
| Medium high constraint | 2.0 (0.37) | 2300 (635) |
| High constraint | 1.1 (0.09) | 2309 (634) |
| Encoding/retrieval task | | |
| | Word classification Mean (s.d.) | Word retrieval Mean (s.d.) |
| Total number correct (out of 36 and 72 stimuli respectively) | 35 (1.3) | 53* (8.2) |
| Total number incorrect | 0.3 (0.6) | 14.2 (5.1) |
| Reaction time | 1037 (440) | 1308 (388) |

* for correct recognition, total correct = 25 (5.0), for correct rejection, total correct = 28 (5.5)

There was also a significant negative correlation between activation in the left posterior lobe of the cerebellum and the suspiciousness/persecution score for sentence completion versus rest ($p = 0.002$), and a more anterior and right sided region of the cerebellum was significantly positively correlated with the delusion score for the parametric contrast ($p = 0.04$), see Figure 4, Table 3.


Experiment 2: Encoding/retrieval task

The key finding from the encoding/retrieval task was a significant negative correlation between the suspiciousness/persecution score and activation in the bilateral MTL for the retrieval task (correct rejection versus experimental baseline activation, right $p = 0.013$, left $p = 0.022$), Figure 3b. There was also a negative correlation with activation in the left anterior and posterior lobes of the cerebellum ($p = 0.052$), Figure 3, Table 3, and a significant positive correlation between activation in the lentiform nucleus and the delusions score ($p = 0.036$).


For word classification versus experimental baseline there was a significant positive correlation between the delusion score and activation in the posterior lobe of the cerebellum ($p = 0.001$) (Figure 4), and a significant negative correlation between activation in the angular gyrus and the hallucinations score ($p = 0.041$), Table 3. There were no significant correlations with correct recognition versus experimental baseline.

Discussion

Although previous studies have found associations between symptoms and functional imaging correlates in patients with established illness, the current study involved a large number of un-medicated high risk subjects experiencing psychotic symptoms and as such avoids potential contamination of findings by anti-psychotic medication and chronicity of disease. We performed these

**Figure 2**

Within group activation maps for sentence completion task (a, b) and verbal encoding/retrieval task (c-e). For the Hayling sentence completion test, contrasts shown are (a) sentence completion versus rest and (b) the parametric contrast representing increasing activation with increasing difficulty. For the encoding/retrieval task, contrasts shown are (c) word classification versus experimental baseline, (d) correct recognition versus experimental baseline, and (e) correct rejection versus experimental baseline. For illustration purposes maps are thresholded at $T = 4.5, 4, 6, 5, 5$ respectively, extent threshold = 50 voxels. Left hemisphere shown on left of image.

**Figure 3**

Correlations with symptom scores in hypothesised regions. (a) significant positive correlation between severity of hallucinations and activation in left middle temporal gyrus for sentence completion task, (b) significant negative correlation between severity of suspiciousness/persecution score and activation in bilateral MTL for the retrieval task. Maps thresholded at $p = 0.005$, $K_E = 300$ voxels. Left hemisphere shown on left of image

analyses by examining the effects of the presence of these symptoms on the neural responses to cognitive tasks. The tasks were not designed specifically to elicit symptoms, but were chosen to involve networks and cognitive processes known to be abnormal in patients with schizophrenia. The results indicated that symptom scores correlated with brain activation in a number of regions broadly consistent with the extant literature. Since in the analysis conducted the three symptom dimensions were modelled simultaneously within one design matrix, intersubject variability common to two or more symptom dimensions is not reflected in the correlation values. The values obtained therefore represent correlations specific to each symptom dimension which is not accounted for by either of the other two. However, due to the approach taken, we would like to state that since all high risk subjects were entered into the analysis, the correlation values may to some extent reflect the presence of symptoms as well as severity. As hypothesised there were significant correla-

Table 3: Correlation between fmri activation and PANSS scores

| Sentence completion task: Sentence completion versus rest | | | | | | | | | |
|--|-----------------------------------|--------------|--------------|---------------------------------|-----------------------------------|--------------|--------------|---|--|
| +ve correlation | | | | | -ve correlation | | | | |
| | P value (Z score;K _E) | Correlation* | Co-ordinates | Region | P value (Z score;K _E) | Correlation* | Co-ordinates | Region | |
| Delusions | n/s | - | - | - | n/s | - | - | - | |
| Hallucinations | 0.01# (4.01; 371) | 0.41 | -48 6 -29 | L middle temporal gyrus (BA 21) | n/s | - | - | - | |
| Suspiciousness/persecution | n/s | - | - | - | 0.002 (4.45; 702) | -0.46 | 0 -74 -35 | L posterior lobe of cerebellum | |
| Sentence completion task: Parametric contrast | | | | | | | | | |
| Delusions | 0.04 (3.83; 337) | 0.40 | 6 -48 -35 | R posterior lobe of cerebellum | n/s | - | - | - | |
| Hallucinations | n/s | - | - | - | n/s | - | - | - | |
| Suspiciousness/persecution | n/s | - | - | - | n/s | - | - | - | |
| Verbal encoding: Word classification versus experimental baseline | | | | | | | | | |
| +ve correlation | | | | | -ve correlation | | | | |
| | P value (Z score;K _E) | Correlation* | Co-ordinates | Region | P value (Z score;K _E) | Correlation* | Co-ordinates | Region | |
| Delusions | 0.001 (4.28; 1097) | 0.45 | 22 -62 -35 | R posterior lobe of cerebellum | n/s | - | - | - | |
| | 0.023 (3.32; 664) | 0.35 | -26 -89 -22 | L posterior lobe of cerebellum | | | | | |
| Hallucinations | n/s | - | - | - | 0.041 (3.51; 582) | -0.38 | 44 -70 27 | R angular gyrus (BA39) | |
| Suspiciousness/persecution | n/s | - | - | - | n/s | - | - | - | |
| Verbal retrieval: Correct recognition versus experimental baseline | | | | | | | | | |
| Delusions | n/s | - | - | - | n/s | - | - | - | |
| Hallucinations | n/s | - | - | - | n/s | - | - | - | |
| Suspiciousness/persecution | n/s | - | - | - | n/s | - | - | - | |
| Verbal retrieval: Correct rejection versus experimental baseline | | | | | | | | | |
| Delusions | 0.036 (3.68; 856) | 0.39 | 22 -12 -6 | R lentiform nucleus | n/s | - | - | - | |
| Hallucinations | n/s | - | - | - | n/s | - | - | - | |
| Suspiciousness/persecution | n/s | - | - | - | 0.013 (3.98; 1083) | -0.42 | 22 -12 -10 | R superior border of hippocampus | |
| | | | | | 0.022 (3.71; 963) | -0.39 | -37 -3 -27 | L lateral border of amygdala/hippocampus | |
| | | | | | 0.052 (3.70; 780) | -0.39 | -7 -28 -41 | L anterior + posterior lobe of cerebellum | |

*determined from partial correlation coefficient analysis in SPSS

#with small volume correction for bilateral temporal lobes




Figure 4
Correlations with symptoms scores in cerebellar regions. Negative correlations with suspiciousness/persecution score for (a) sentence completion, (b) verbal retrieval. Positive correlations with delusions score for (c) sentence completion, (d) verbal encoding. Maps thresholded at $p = 0.005$ uncorrected, extent threshold = 100 voxels. Left hemisphere shown on left of image

tions with the hallucination score and activation in the left lateral temporal cortex for the sentence completion task. For the encoding/retrieval task there were significant correlations with the suspiciousness/persecution score and activation in the bilateral MTL. Across both tasks there were significant correlations between delusions and suspiciousness/persecution scores and activation in the cerebellum, with different patterns of laterality.

The finding of an association between the hallucinations measure and activation in lateral temporal cortex is consistent with other reports linking this region to auditory hallucinations [9-11,14], and with cognitive neuropsychological theories of auditory verbal hallucinations regarding the misattribution of inner speech [1,34]. Although regions most typically associated with hallucinations are mid to posterior sections of the superior temporal gyrus, other studies have indeed reported associations with activation in the middle temporal gyrus [10,14,16]. The cognitive model most consistent with our task and results is that auditory hallucinations represent sub-vocal verbalisation which is not associated with the usual inhibition of auditory cortex that signifies self-production [1].

The memory task revealed strong negative correlations between the suspiciousness/persecution score and activation in bilateral MTL structures. This association was seen for correct rejection versus experimental baseline and not correct recognition. There is previous evidence however to indicate MTL regions are preferentially involved in correctly identifying new words (correct rejection) compared to correctly identifying old words (correct recognition) [35]. Previous literature has also implicated MTL regions in the formation of positive psychotic symptoms [3-5,36], and those that have examined delusional beliefs specifically also report decreased activation with increasing symptom severity [36]. There is also substantial evidence for structural abnormalities in this region in patients with established schizophrenia and high risk individuals [37-39]. The MTL, including the amygdala and hippocampus, are known to be involved in emotional, mnemonic and social processing [40-42] and dysregulation of these processes may contribute to the development of persecutory beliefs. For example, persecutory beliefs have been hypothesized to arise from a disruption in the processes mediating the formation and maintenance of normal social beliefs, and patients with persecutory beliefs have been shown to attend excessively to threat-related stimuli and to preferentially recall such information [43]. Impaired processing of threat-related stimuli has also been previously linked to dysfunction of the amygdala in paranoid patients with schizophrenia [44].

Although not part of our original hypothesis, we also found correlations between both suspiciousness/persecution and delusion scores and activation of the cerebellum across both of the tasks. Abnormal cerebellar function has been previously reported in the literature on the established illness [45] and in those at high genetic risk [26]. Further, it has been reported that patients with posterior lobe lesions have deficits which are not restricted to motor function, but also present affective disturbances, disinhibition, and psychotic symptoms [46,47]. The cerebellum has also been implicated in the 'cognitive dysmetria' model of schizophrenia where dysfunction in cortico-cerebellar-thalamic-cortical circuits are proposed to result in the abnormal synchrony or co-ordination of mental processing, ultimately resulting in the clinical features of the illness [48]. Although there are few imaging studies of patients with persecutory delusions, neuropsychological studies have repeatedly demonstrated that patients commonly attribute negative events to external sources [49]. Neurobiological theories have also directly implicated the cerebellum regarding deficits of 'internal monitoring' which are suggested, for example, to underlie the formation of delusions of alien control where the patient attributes their own actions to an external agent [19]. The current findings are therefore consistent with a growing

literature implicating a role for the cerebellum in the formation of delusions.

The nature of the correlations in the cerebellum are however complex. The correlations are primarily right sided with regards to the delusions score, and primarily left sided with the suspiciousness/persecution score. It has been suggested that different types of delusions may differ pathophysiologically and there is an important role for emotion in the formation of paranoid/suspicious delusions via their effects on memory [49]. Following the model of brain function where emotion is predominantly associated with right sided cerebral regions, crossed cerebellar connections could mean that left sided cerebellar regions would be more involved in emotional/affective components of delusions, consistent with the present findings. Furthermore, since emotion is considered to be involved via its effect on memory processes, our MTL associations with the severity of suspiciousness/persecution also fits with this theory. We however recognise the speculative nature of these findings, particularly with regard to the paucity of other studies examining delusional beliefs. Further, in the group we have examined many had both hallucinations and delusions (including persecutory and non-persecutory) and it is therefore difficult to disambiguate individual components. That said, however, the fact that this was a study examining correlations with symptom scores reduces the chances of confounding effects by other symptoms. Studies directly manipulating experimental affective components in subjects with isolated symptoms may help clarify these results.

Conclusion

In conclusion, these results support the view that hallucinations and delusions have differential pathophysologies, that the lateral temporal cortex is associated with the experience of hallucinations, and that the MTL is associated with the production of, or perhaps the response to, suspiciousness/persecutory beliefs. The more novel finding of the involvement of cerebellar structures was seen across both tasks with both delusional and suspiciousness/persecution scores with differing patterns of laterality. That the current results are seen in un-medicated high risk subjects indicates these associations are not specific to the established illness and are not related to medication effects and hence may have relevance for the understanding of the biological basis of the disorder.

Competing interests

The author(s) declare that they have no competing interests.

Authors' contributions

HCW, VEG, JH, ES and DEJ contributed to analysis of the functional imaging data. AM contributed to statistical

analysis. DGCO, ECJ and SML conducted clinical evaluations, conceived the original study, and participated in its design and co-ordination. All authors contributed to drafting of the manuscript and have read and approved the final version.

Acknowledgements

The study was funded by an MRC programme grant. Dr Lawrie and Dr Whalley are supported by a charitable donation from the Dr Mortimer and Theresa Sackler Foundation. Scanning was carried out at the Brain Imaging Research Centre (BIRC) for Scotland. Both BIRC and the Centre for Functional Imaging studies (CFIS) are funded by the Scottish Higher Education Funding Council (SHEFC). We would also like to thank those involved in subject recruitment, the radiographers, and the study participants.

References

1. Frith: **The Cognitive Neuropsychology of Schizophrenia**. Hove, Lawrence Erlbaum Associates; 1992.
2. Weiss AP, Heckers S: **Neuroimaging of hallucinations: a review of the literature**. *Psychiatry Res* 1999, **92**(2-3):61-74.
3. Liddle PF, Friston KJ, Frith CD, Hirsch SR, Jones T, Frackowiak RS: **Patterns of cerebral blood flow in schizophrenia**. *Br J Psychiatry* 1992, **160**:179-186.
4. Friston KJ, Liddle PF, Frith CD, Hirsch SR, Frackowiak RS: **The left medial temporal region and schizophrenia. A PET study**. *Brain* 1992, **115**(Pt 2):367-382.
5. Surguladze S, Russell T, Kucharska-Pietura K, Travis MJ, Giampietro V, David AS, Phillips ML: **A reversal of the normal pattern of parahippocampal response to neutral and fearful faces is associated with reality distortion in schizophrenia**. *Biol Psychiatry* 2006, **60**(5):423-431.
6. Honey GD, Sharma T, Suckling J, Giampietro V, Soni W, Williams SC, Bullmore ET: **The functional neuroanatomy of schizophrenic subsyndromes**. *Psychol Med* 2003, **33**(6):1007-1018.
7. Menon V, Agnoston RT, Mathalon DH, Glover GH, Pfefferbaum A: **Functional neuroanatomy of auditory working memory in schizophrenia: relation to positive and negative symptoms**. *Neuroimage* 2001, **13**(3):433-446.
8. Yuasa S, Kurachi M, Suzuki M, Kadono Y, Matsui M, Saitoh O, Seto H: **Clinical symptoms and regional cerebral blood flow in schizophrenia**. *Eur Arch Psychiatry Clin Neurosci* 1995, **246**(1):7-12.
9. David AS, Woodruff PW, Howard R, Mellers JD, Brammer M, Bullmore E, Wright I, Andrew C, Williams SC: **Auditory hallucinations inhibit exogenous activation of auditory association cortex**. *Neuroreport* 1996, **7**(4):932-936.
10. Dierks T, Linden DE, Jandl M, Formisano E, Goebel R, Lanfermann H, Singer W: **Activation of Heschl's gyrus during auditory hallucinations**. *Neuron* 1999, **22**(3):615-621.
11. Lennox BR, Park SB, Medley I, Morris PG, Jones PB: **The functional anatomy of auditory hallucinations in schizophrenia**. *Psychiatry Res* 2000, **100**(1):13-20.
12. McGuire PK, Shah GM, Murray RM: **Increased blood flow in Broca's area during auditory hallucinations in schizophrenia**. *Lancet* 1993, **342**(8873):703-706.
13. Shergill SS, Brammer MJ, Fukuda R, Williams SC, Murray RM, McGuire PK: **Engagement of brain areas implicated in processing inner speech in people with auditory hallucinations**. *Br J Psychiatry* 2003, **182**:525-531.
14. Shergill SS, Brammer MJ, Williams SC, Murray RM, McGuire PK: **Mapping auditory hallucinations in schizophrenia using functional magnetic resonance imaging**. *Arch Gen Psychiatry* 2000, **57**(11):1033-1038.
15. Woodruff PW, Wright IC, Bullmore ET, Brammer M, Howard RJ, Williams SC, Shapleske J, Rossell S, David AS, McGuire PK, Murray RM: **Auditory hallucinations and the temporal cortical response to speech in schizophrenia: a functional magnetic resonance imaging study**. *Am J Psychiatry* 1997, **154**(12):1676-1682.
16. Silbersweig D, Stern E: **Functional neuroimaging of hallucinations in schizophrenia: toward an integration of bottom-up and top-down approaches**. *Mol Psychiatry* 1996, **1**(5):367-375.

17. Cleghorn JM, Franco S, Szechtman B, Kaplan RD, Szechtman H, Brown GM, Nahmias C, Garnett ES: **Toward a brain map of auditory hallucinations.**[see comment]. *Am J Psychiatry* 1992, **149**(8):1062-1069.
18. Blackwood NJ, Bentall RP, Fyftche DH, Simmons A, Murray RM, Howard RJ: **Persecutory delusions and the determination of self-relevance: an fMRI investigation.**[see comment]. *Psychol Med* 2004, **34**(4):591-596.
19. Blakemore SJ, Oakley DA, Frith CD: **Delusions of alien control in the normal brain.** *Neuropsychologia* 2003, **41**(8):1058-1067.
20. Spence SA, Brooks DJ, Hirsch SR, Liddle PF, Meehan J, Grasby PM: **A PET study of voluntary movement in schizophrenic patients experiencing passivity phenomena (delusions of alien control).** *Brain* 1997, **120**(Pt 11):1997-2011.
21. Verdoux H, van Os J: **Psychotic symptoms in non-clinical populations and the continuum of psychosis.** *Schizophr Res* 2002, **54**(1-2):59-65.
22. Johnstone EC, Abukmeil SS, Byrne M, Clafferty R, Grant E, Hodges A, Lawrie SM, Owens DG: **Edinburgh high risk study--findings after four years: demographic, attainment and psychopathological issues.** *Schizophr Res* 2000, **46**(1):1-15.
23. Hemsley DR: **The development of a cognitive model of schizophrenia: placing it in context.** *Neurosci Biobehav Rev* 2005, **29**(6):977-988.
24. Hodges A, Byrne M, Grant E, Johnstone E: **People at risk of schizophrenia. Sample characteristics of the first 100 cases in the Edinburgh High-Risk Study.**[see comment]. *Br J Psychiatry* 1999, **174**:547-553.
25. Johnstone EC, Ebmeier KP, Miller P, Owens DG, Lawrie SM: **Predicting schizophrenia: findings from the Edinburgh High-Risk Study.** *Br J Psychiatry* 2005, **186**:18-25.
26. Whalley HC, Simonotto E, Flett S, Marshall I, Ebmeier KP, Owens DG, Goddard NH, Johnstone EC, Lawrie SM: **fMRI correlates of state and trait effects in subjects at genetically enhanced risk of schizophrenia.**[see comment]. *Brain* 2004, **127**(Pt 3):478-490.
27. Whyte MC, Whalley HC, Simonotto E, Flett S, Shillcock R, Marshall I, Goddard NH, Johnstone EC, Lawrie SM: **Event-related fMRI of word classification and successful word recognition in subjects at genetically enhanced risk of schizophrenia.** *Psychol Med* 2006, **36**(10):1427-1439.
28. Kay SR, Fiszbein A, Opler LA: **The positive and negative syndrome scale (PANSS) for schizophrenia.** *Schizophr Bull* 1987, **13**(2):261-276.
29. Owens DG, Finlayson A, Mercer G, Johnstone EC: **The use of videotaped assessment in relation to a study of enhanced management in treatment-resistant schizophrenia.** *J Psychopharmacol (Oxf)* 1997, **11**(4):357-360.
30. Bloom PA, Fischler I: **Completion norms for 329 sentence contexts.** *Mem Cognit* 1980, **8**(6):631-642.
31. **SPM2, The Wellcome Department of Cognitive Neurology and collaborators, Institute of Neurology, London** [<http://www.fil.ion.ucl.ac.uk/spm/>]
32. Whalley HC, Kestelman JN, Rimmington JE, Kelso A, Abukmeil SS, Best JJ, Johnstone EC, Lawrie SM: **Methodological issues in volumetric magnetic resonance imaging of the brain in the Edinburgh High Risk Project.** *Psychiatry Res* 1999, **91**(1):31-44.
33. **CBU Imaging** [<http://www.mrc-cbu.cam.ac.uk/Imaging>]
34. Ford JM, Mathalon DH: **Corollary discharge dysfunction in schizophrenia: can it explain auditory hallucinations?** *Int J Psychophysiol* 2005, **58**(2-3):179-189.
35. Henson RN, Hornberger M, Rugg MD: **Further dissociating the processes involved in recognition memory: an fMRI study.** *J Cogn Neurosci* 2005, **17**(7):1058-1073.
36. Lahti AC, Weiler MA, Holcomb HH, Tamminga CA, Carpenter WT, McMahon R: **Correlations between rCBF and symptoms in two independent cohorts of drug-free patients with schizophrenia.** *Neuropsychopharmacology* 2006, **31**(1):221-230.
37. Job DE, Whalley HC, McConnell S, Glabus M, Johnstone EC, Lawrie SM: **Voxel-based morphometry of grey matter densities in subjects at high risk of schizophrenia.** *Schizophr Res* 2003, **64**(1):1-13.
38. Lawrie SM, Abukmeil SS: **Brain abnormality in schizophrenia. A systematic and quantitative review of volumetric magnetic resonance imaging studies.**[see comment]. *Br J Psychiatry* 1998, **172**:110-120.
39. Wright IC, McGuire PK, Poline JB, Travers JM, Murray RM, Frith CD, Frackowiak RS, Friston KJ: **A voxel-based method for the statistical analysis of gray and white matter density applied to schizophrenia.** *Neuroimage* 1995, **2**(4):244-252.
40. Adolphs R: **The neurobiology of social cognition.** *Curr Opin Neurol* 2001, **11**(2):231-239.
41. LaBar KS, Cabeza R: **Cognitive neuroscience of emotional memory.** *Nature Reviews Neuroscience* 2006, **7**(1):54-64.
42. Phelps EA, LeDoux JE: **Contributions of the amygdala to emotion processing: from animal models to human behavior.** *Neuron* 2005, **48**(2):175-187.
43. Blackwood NJ, Howard RJ, Bentall RP, Murray RM: **Cognitive neuropsychiatric models of persecutory delusions.** *Am J Psychiatry* 2001, **158**(4):527-539.
44. Williams LM, Das P, Harris AW, Liddell BB, Brammer MJ, Olivieri G, Skerrett D, Phillips ML, David AS, Peduto A, Gordon E: **Dysregulation of arousal and amygdala-prefrontal systems in paranoid schizophrenia.** *Am J Psychiatry* 2004, **161**(3):480-489.
45. Ho BC, Mola C, Andreasen NC: **Cerebellar dysfunction in neuroleptic naive schizophrenia patients: clinical, cognitive, and neuroanatomic correlates of cerebellar neurologic signs.** *Biol Psychiatry* 2004, **55**(12):1146-1153.
46. Schmahmann JD: **The role of the cerebellum in affect and psychosis.** *Journal of Neurolinguistics* 2000, **13**:189-214.
47. Schmahmann JD, Sherman JC: **The cerebellar cognitive affective syndrome.**[see comment]. *Brain* 1998, **121**(Pt 4):561-579.
48. Andreasen NC, O'Leary DS, Cizadlo T, Arndt S, Rezaei K, Boles Ponto LL, Watkins GL, Hichwa RD: **Schizophrenia and cognitive dysmetria: a positron-emission tomography study of dysfunctional prefrontal-thalamic-cerebellar circuitry.** *Proceedings of the National Academy of Sciences of the United States of America* 1996, **93**:9985-9990.
49. Gilleen J, David AS: **The cognitive neuropsychiatry of delusions: from psychopathology to neuropsychology and back again.** *Psychol Med* 2005, **35**(1):5-12.

Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1471-244X/7/61/prepub>

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

